

Brush Busters Soil Spot Spray Method

Works Best: On blueberry or redberry cedars less than 3 feet tall.

When to Apply: Late winter to mid-spring (ideally, before expected rainfall).

① Prepare Equipment

Soil spot sprays should be applied with an exact delivery handgun. This equipment is available from most herbicide retail outlets. The handgun delivers a thin stream of predetermined volume when triggered. Adjust the handgun to deliver 2 ml (cc) for each pull of the trigger. If only a few plants are to be treated, a disposable syringe can be used. All spray equipment should be thoroughly cleaned immediately after use.

② Prepare Herbicide

Velpar L™ is the recommended herbicide for soil spot sprays to control cedar. The herbicide is used undiluted, by attaching an exact delivery handgun or syringe to the herbicide container.

③ Apply the Herbicide

Apply undiluted Velpar L™ to the soil surface midway between the cedar stem and the canopy edge. Apply 2 ml for every 3 feet of plant height or every 3 feet of plant canopy diameter

(whichever is greater). If plant size requires more than a single 2-ml application, space applications equally around the plant. Apply each 2-ml dose to a single spot on the soil surface. On slopes, apply most of the herbicide on the uphill side of the stem.



Keep these points in mind:

- Follow herbicide label directions.
- Do not use on marshy or poorly drained sites.
- Do not use on clay soils.
- Do not apply to snow-covered or frozen ground.
- Do not apply within three times the height or canopy diameter (whichever is greater) of desirable trees such as oaks or pecans.
- Rainfall is required to "activate" Velpar L™ in the soil. Plants will begin to show symptoms within 3 to 6 weeks of initial rainfall. One to three growing seasons may be required before plants die, and during this time cedars usually sprout new leaves several times.
- Grasses and weeds will be killed where each spot of Velpar L™ is applied. Recovery may take 2 to 3 years.
- Cost of treatment escalates rapidly as density and size of cedar increases.

Brush Busters Top Removal Method

Blueberry cedar: Remove the plant top at or near the ground line with a chain saw, pruner, axe, etc. Seedlings (plants less than 2 feet tall) can be killed easily by hand grubbing.

Redberry cedar: Since redberry cedar resprouts, top removal will not effectively control plants that are over 8 to 10 years old. Redberry cedar seedlings (plants less than 2 feet tall) can be easily killed if they are grubbed below the basal "knob," located at or slightly beneath the soil surface.



Texas Agricultural Extension Service
Texas Agricultural Experiment Station
The Texas A&M University System



How to Master Cedar

Three safe and effective ways to control small cedar (juniper).

Individual Plant Treatment Series

Allan McGinty, Professor and Extension Range Specialist,
The Texas A&M University System
Darrell Ueckert, Professor, Rangeland Ecology
and Management, Texas Agricultural Experiment Station,
The Texas A&M University System

Educational programs of the Texas Agricultural Extension Service are open to all people without regard to race, color, sex, disability, religion, age or national origin.

Issued in furtherance of Cooperative Extension Work in Agriculture and Home Economics, Acts of Congress of May 8, 1914, as amended, and June 30, 1914, in cooperation with the United States Department of Agriculture. Zerle L. Carpenter, Director, Texas Agricultural Extension Service, The Texas A&M University System.
11M, 6-97, Revised

RS

Texas Agricultural Extension Service, Zerle L. Carpenter, Director
Texas Agricultural Experiment Station, Edward A. Hilger, Director
The Texas A&M University System, College Station, Texas

Brush Busters Leaf Spray Method

Cedars are very aggressive and damaging woody plants on Texas rangelands. There are two major cedar species west of Interstate Highway 35, redberry cedar and blueberry cedar. It is important to know which species you have because the treatments vary. As the names imply, blueberry cedar has blue berries and redberry cedar has red berries. In addition, redberry cedar has small specks of white wax on its leaves and twigs. Blueberry cedar is more common in central and south-central Texas, while redberry cedar is more common in west, west-central and north-central Texas. It is not uncommon for the two species to occur together.

Controlling cedar is not a one-time job. Livestock and wildlife spread the seeds easily, so you'll need to check your land occasionally and treat unwanted seedlings. It is very important to control cedar as seedlings and saplings. Controlling mature cedars requires heavy equipment or large quantities of herbicides and is usually very expensive. Using these Brush Buster methods, you'll be able to keep the plants you want and get rid of those you don't.

Brush Busters recommends three ways to control cedar. Two herbicide treatments—leaf spray and soil spot spray—work best on cedar less than 3 feet tall. The top removal method, which involves cutting the tree at ground level, will control blueberry cedar, but redberry cedar must be grubbed (cut) below the soil surface. These three “select” methods are easy, inexpensive, environmentally responsible, and effective. Your results may vary with the weather and other conditions, but you should be able to knock out more than 7 of 10 cedars treated.

Works Best: On blueberry or redberry cedars less than 3 feet tall.

When to Apply: Spring through summer, when cedar is actively growing.

① Prepare Equipment

Small pump-up garden sprayers, backpack sprays, cattle sprayers, or sprayers mounted on 4-wheel-drive all-terrain vehicles (ATVs) work well. Garden sprayers are best for treating a few plants, and backpack sprayers are usually more efficient for larger acreages or higher cedar densities. ATV sprayers become more efficient on very large acreages or as the distance between plants increases. Make sure your spraygun has an adjustable cone nozzle (X6 to X8 orifice size) capable of delivering a coarse spray (large droplets).

② Mix Herbicide Spray


You can achieve 76 to 100 percent rootkill of small cedar by spraying with the herbicide Tordon 22K™. Prepare the spray mix as a 1% concentration of Tordon 22K™ in water (see table). To ensure a thorough coating of the cedar leaves,

Recommended leaf spray for cedar.*

Ingredient	Concentration in spray solution	Tank size		
		3-gal.	14-gal.	25-gal.
Tordon 22K™	1%	4 oz.	18 oz.	32 oz.
Surfactant or dishwashing detergent	1/4%	1 oz.	5 oz.	8 oz.
Hi-Light Blue Dye™	1/4-1/2%	1-2 oz.	5-9 oz.	8-16 oz.

*All spray solutions are mixed in water.

add commercial surfactant or liquid dishwashing detergent to the spray mix. It may be helpful to add a dye, such as Hi-Light Blue Dye™, to mark plants that have been sprayed.



③ Spray the Cedar

Thoroughly wet all the leaves of each cedar plant to the point of runoff.

Keep these points in mind:

- The purchase of Tordon 22K™ requires a Pesticide Applicator License from the Texas Department of Agriculture. See your county Extension agent for license information.
- Follow herbicide label directions.
- Do not spray within 100 feet of known sinkholes or fractures that would allow herbicide to enter underground water aquifers.
- Do not spray if cedar foliage is wet.
- Do not spray immediately upwind of desirable trees, shrubs or susceptible crops.
- Cost of treatment increases rapidly as density and size of cedar increases.
- Large, mature blueberry cedar may be nesting habitat for the endangered golden-cheeked warbler. If in doubt, contact the U.S. Fish & Wildlife Service before treating.
- Do not treat large numbers of cedars beneath the canopies or within three times the height or diameter of desirable trees such as oaks or pecans.

The information given herein is for educational purposes only. Reference to commercial products or trade names is made with the understanding that no discrimination is intended and no endorsement by the Cooperative Extension Service or Texas Agricultural Experiment Station is implied.